



Contribution ID: 163

Type: Poster

Kinetics of neptunium(V) conversion in strong nitric acid solutions containing potassium phosphotungstate, $K_{10}P_2W_{17}O_{61}$

Monday, 19 April 2010 11:45 (20 minutes)

Behavior of Np(V) in strong nitric acid solutions with different strength (1,0 ÷ 3,0) mol·l⁻¹ and KPW (1 ÷ 5)·10⁻³ mol·l⁻¹, containing potassium phosphotungstate, K₁₀P₂W₁₇O₆₁ (KPW) is examined by spectrophotometric method .

It is established that Np (V) final conversion products under studied experimental conditions are Np (IV) and Np (VI), and the process is going in accordance with a first-order rate law in regard to neptunium(V) concentration.

Primary author: Dr CHISTYAKOV, Vladimir (Russia, Rosatom, JSC"State Scientific Center-Research Institute of Atomic Reactors", Radiochemical Unit)

Presenter: Dr CHISTYAKOV, Vladimir (Russia, Rosatom, JSC"State Scientific Center-Research Institute of Atomic Reactors", Radiochemical Unit)

Session Classification: Poster Session - Chemistry of Actinide and Trans-actinide Elements

Track Classification: Chemistry of Actinide and Trans-actinide Elements