



Contribution ID: 225

Type: Verbal

NAA for studying detoxification of Cr and Hg by *Arthrobacter globiformis*

Bacterial reduction and detoxification of potentially toxic metals is one of most promising strategies for the bioremediation of contaminated environmental media. In our previous studies we have established that the Gram-positive bacterial strain *Arthrobacter globiformis* isolated from basalts taken from ecologically mostly polluted region of Georgia (Marneuli) can reduce and detoxify of Cr(VI) with high efficiency [1]. In the present investigation instrumental neutron activation analysis (INAA) was applied to study (1) accumulation of Cr(VI) in *A. globiformis* in the presence of Hg(II); (2) accumulation of Hg(II) in bacterial cells; and (3) effects of Hg(II) and mixture of Cr(VI) - Hg(II) on the elemental composition of bacteria. Our experiments were focused on the dose-dependent effects of Cr(VI) and Hg(II). Cr(VI) as $[K_2CrO_4]$ and Hg(II) as $[Hg(NO_3)_2 \cdot H_2O]$ were added to the bacterial cell cultures at an early stationary phase of their growth. Two sets of experiments were performed. In the first set the concentration of Hg(II) varied within the range of 50–5000 $\mu\text{g/L}$. In the second set a 500 $\mu\text{g/L}$ concentration of Hg(II) was added to the bacterial cells at each given concentration of Cr(VI) within the range of 50–1000 mg/L . According to the results obtained, the dose-dependent character of Cr(VI) accumulation by the tested bacterial strain was not significantly affected by the presence of Hg(II). Accumulation of Hg(II), similar to the Cr(VI) accumulation, follows well the Langmuir-Freundlich model. Besides, NAA measurements showed increased content of Fe in bacteria under Hg and Cr action, suggesting that Fe-containing biomolecules play a decisive role in detoxifying of heavy metals by *A. globiformis*. A concentration of 5000 $\mu\text{g/L}$ of Hg(II) was found to be critical for *A. globiformis*. At this concentration of Hg(II) the concentrations of both essential (Na, Mg, Al, Cl, K, Mn, Zn) and some non-essential elements (Rb, Sb, Sc, As) changed drastically along with a decrease of the biomass of bacteria by a factor of 2. One may assume that under this high exposure to Hg(II) the structure of the bacterial cell wall was destroyed. Acknowledgement: We are grateful to STCU for their support (Grant #4330)

Reference: N. Tsibakhashvili, L. Mosulishvili, T. Kalabegishvili, E. Kirkesali, I. Murusidze, S. Kerkenjia, M. Frontasyeva, H.-Y. Holman, "Biotechnology of Cr(VI) transformation into Cr(III) complexes", *J. Radioanal. Nucl. Chem.* (2008), 278(3), 565-569.

Keywords *Arthrobacter globiformis*, NAA, Cr(VI), Hg(II), accumulation

Primary author: Dr TSIBAKHASHVILI, Nelly (Andronikashvili Institute of Physics, Tbilisi, Georgia)

Co-authors: Dr KIRKESALI, E. (Andronikashvili Institute of Physics, Tbilisi, Georgia); Dr MURUSIDZE, I. (Chavchavadze State University, Tbilisi 0179, Georgia); Ms ZINICOVSCAIA, I. (Joint Institute for Nuclear Research); Dr MOSULISHVILI, L. (Andronikashvili Institute of Physics, Tbilisi, Georgia); Dr FRONTASYEVA, M.V. (Joint Institute for Nuclear Research); Dr BODE, P. (Delft University of Technology, 2629JB Delft, The Netherlands); Dr TH.G.VAN MEERTEN, Th. (Delft University of Technology, 2629JB Delft, The Netherlands)

Presenter: Dr FRONTASYEVA, M.V. (Joint Institute for Nuclear Research)

Track Classification: Nuclear Analytical Methods