



Contribution ID: 200

Type: Verbal

## **The Chernobyl-derived $^{137}\text{Cs}$ accumulation in artificial reservoirs in Central Russia as indicator of sediment sources**

*Tuesday, 10 September 2019 16:30 (15 minutes)*

**Primary authors:** Mr IVANOV, Maksim (Faculty of Geography, Lomonosov Moscow State University, Moscow, 119991, Russia; Institute of Geography, Russian Academy of Science, Moscow, 119017, Russia); Mr KONSTANTINOV, Evgenyi (Institute of Geography, Russian Academy of Science, Moscow, 119017, Russia); Mrs IVANOVA, Nadezhda (Faculty of Geography, Lomonosov Moscow State University, Moscow, 119991, Russia); Mr TSYPLENKOV, Anatoly (Faculty of Geography, Lomonosov Moscow State University, Moscow, 119991, Russia); Mrs KUZMENKOVA, Natalya (Faculty of Chemistry, Lomonosov Moscow State University, Moscow, 119991, Russia); Mr GURINOV, Artem (Faculty of Geography, Lomonosov Moscow State University, Moscow, 119991, Russia); Prof. KONOPLEV, Alexei (Institute of Environmental Radioactivity, Fukushima University, Fukushima, 960-1296, Japan); Dr GOLOSOV, Valentin (Faculty of Geography, Lomonosov Moscow State University, Moscow, 119991, Russia; Institute of Geography, Russian Academy of Science, Moscow, 119017, Russia)

**Presenter:** Mr IVANOV, Maksim (Faculty of Geography, Lomonosov Moscow State University, Moscow, 119991, Russia; Institute of Geography, Russian Academy of Science, Moscow, 119017, Russia)

**Session Classification:** Parallel Session 5A