



Contribution ID: 1041

Type: Verbal

EVALUATION OF THE DEPENDENCE OF THE ACTIVITY CONCENTRATIONS RATIOS $^{234}\text{U}/^{238}\text{U}$ AND $^{235}\text{U}/^{238}\text{U}$ IN WATERS ON THE ^{238}U ACTIVITY CONCENTRATION

Wednesday, 18 May 2022 09:24 (18 minutes)

In routine radiochemical analyzes of water, especially drinking water, in recent years we have analyzed more than 4,000 water samples for uranium ^{238}U , ^{235}U and ^{234}U isotope contents by ICP/SFMS method in the ALS Scandinavia AB laboratory, Luleå, Sweden and the Central Analytical Laboratory of ÚJV Řež, a.s., Husinec – Řež, in the Czech Republic. By statistical processing of an extensive set of data, we evaluated the interesting patterns that are discussed in this article and which can significantly help in the evaluation and interpretation of measurements of natural radionuclides in drinking water.

While the $^{235}\text{U}/^{238}\text{U}$ activity ratios in natural waters and rocks are more or less constant, the $^{234}\text{U}/^{238}\text{U}$ activity ratios in natural waters, especially groundwater, but also in some soils, sediments and rocks, usually do not correspond to the calculated theoretical values. The $^{234}\text{U}/^{238}\text{U}$ activity ratio is 1.0 only in closed systems, e.g. in some minerals that have not been subjected to any chemical effect for a long time. In waters, the activity ratio of $^{234}\text{U}/^{238}\text{U}$ is often many times higher than 1.0. The reason for this is the fact that ^{234}U in rocks is formed by the decay of its parent nuclide, ^{238}U . After one alpha decay and two beta decays, the daughter ^{234}U is released from its original position in the crystal lattice of the mineral due to “atomic recoil”, and upon subsequent chemical attack of the rock by groundwater, ^{234}U atoms enter the aquatic environment much more easily than its parent ^{238}U atoms. Thus, uranium-234 isotope is much more chemically reactive than uranium-238 in the solid sample, and as a result, the $^{234}\text{U}/^{238}\text{U}$ activity ratio in waters and in the formed secondary minerals and rocks is much higher than 1.0. The fact that ^{234}Th , the short-term alpha decay product of ^{238}U , can be released to a greater extent into the aqueous environment after atomic recoil, where it decays to ^{234}U very quickly, also contributes in part to the increase in ^{234}U activity against ^{238}U .

We confirmed these facts by statistical processing of an extensive data set of ^{238}U , ^{235}U and ^{234}U contents in the analyzed water samples. The $^{235}\text{U}/^{238}\text{U}$ activity ratio is constant within the uncertainties of the determination, no exception was found. In contrast, the activity ratio of $^{234}\text{U}/^{238}\text{U}$ is significantly higher than 1.0, the overall average of this ratio is 1.9. The value of this ratio depends on the hydro-chemical conditions of the environment, but it can generally be said that the lower the total volume concentration of uranium resp. the volume activity of ^{238}U in groundwater, the higher the value of this ratio. If the volume activity of ^{238}U in groundwater decreases from XX-XXX Bq/L to 0.0XX Bq/L, the 95% percentile distribution of the $^{234}\text{U}/^{238}\text{U}$ ratio will increase from about 3 to 8-10.

The work also discusses relatively “rare cases” where the activity ratio $^{234}\text{U}/^{238}\text{U}$ is less than 1.0.

The values of these activity ratios were also evaluated for about 500 analyzed solid samples.

Primary author: BOUDA, Tomáš (ALS Czech Republic, s.r.o.)

Presenter: BOUDA, Tomáš (ALS Czech Republic, s.r.o.)

Session Classification: Environmental Radioactivity

Track Classification: Radionuclides in the Environment, Radioecology