Workshop on Modern Trends in Quantum Theory



Contribution ID: 13

Type: Talk

Towards Morse theory of dispersion relations

Tuesday, 24 May 2022 15:30 (1 hour)

The question of optimizing an eigenvalue of a family of self-adjoint operators that depends on a set of parameters arises in diverse areas of mathematical physics. Among the particular motivations for this talk are the Floquet-Bloch decomposition of the Schroedinger operator on a periodic structure, nodal count statistics of eigenfunctions of quantum graphs, conical points in potential energy surfaces in quantum chemistry and the minimal spectral partitions of domains. In each of these problems one seeks to identify and/or count the critical points

of the eigenvalue with a given label (say, the third lowest) over the parameter space which is often known and simple, such as a torus.

Classical Morse theory is a set of tools connecting the number of critical points of a smooth function on a manifold to the topological invariants of this manifold. However, the eigenvalues are not smooth due to presence of eigenvalue multiplicities or diabolical points". We rectify this problem for eigenvalues of generic families of finite-dimensional operators. The correctMorse indices" of the problematic diabolical points turn out to be universal: they depend only on the total multiplicity at the diabolical point and on the relative position of the eigenvalue of interest in the eigenvalue group.

Based on a joint work with I.Zelenko.

Primary author: Prof. BERKOLAIKO, Gregory (Texas A&M University, College Station, TX.)

Presenter: Prof. BERKOLAIKO, Gregory (Texas A&M University, College Station, TX.)

Session Classification: Talk