



Contribution ID: 339

Type: Verbal

Actinide stripping by a hydrophilic BTP ligand in aqueous HNO₃ from TODGA-containing organic phase

Monday, 12 May 2014 16:30 (15 minutes)

N,N,N',N'-tetraoctyl-diglycolamide (TODGA) is broadly studied as a non-specific extractant for actinides and lanthanides from HNO₃ solutions of nuclear waste, in particular for group actinide extraction (GANEX process). Geist and his coworkers proposed a novel hydrophilic ligand, 2,6-bis(5,6-di-(sulfophenyl)-1,2,4-triazin-3-yl)pyridine (SO₃-Ph-BTP) for selective stripping of actinides(III) from the loaded organic phase [A. Geist et al., Solvent Extr. Ion Exch., 30, 433 (2012)], and determined (by TRLFS) the stability constants of its consecutive (from 1:1 to 1:3) complexes with Cm(III) and Eu(III) in aqueous solutions [C.M. Ruff et al., Dalton Trans., 41, 14594 (2012)].

In the present work, we studied the distribution of some actinides(III-VI) in the two-phase systems: TODGA (in kerosene) / SO₃-Ph-BTP (0–100 mM in aqueous HNO₃); and determined the apparent stability constants of their complexes in the aqueous phase, based on the distribution ratios of the metal ions. The effect of protonation of the SO₃-Ph-BTP ligand in the acidic solutions was discussed.

This work was financed from the National Centre for Research and Development through the Strategic Program Technologies Supporting Development of Safe Nuclear Power Engineering, task 4: Development of spent nuclear fuel and radioactive waste management techniques and technologies.

Primary author: Mr STECZEK, Łukasz (Institute of Nuclear Chemistry and Technology)

Co-authors: Mr NARBUTT, Jerzy (Institute of Nuclear Chemistry and Technology); Mrs REJNIS, Magdalena (Institute of Nuclear Chemistry and Technology)

Presenter: Mr STECZEK, Łukasz (Institute of Nuclear Chemistry and Technology)

Session Classification: Separation Methods, Speciation 1

Track Classification: Separation Methods, Speciation