



Contribution ID: 71

Type: Poster

Theoretical consideration of the specific activity of Tc-99m produced by the Mo-100(p,2n)Tc-99m reaction at cyclotrons

Thursday, 15 May 2014 17:30 (1h 15m)

Even today the generator-produced radionuclide ^{99m}Tc is used for the vast majority of nuclear medical diagnoses. The parent radionuclide ^{99}Mo is generally produced via fission of highly-enriched ^{235}U , however, at a few nuclear reactor sites in the world only. Due to ageing reactors the world supply of fission-produced ^{99}Mo has become somewhat insecure over the last few years. Alternative methods of direct production of ^{99m}Tc using accelerators are therefore presently attracting great attention. The (p,2n) reaction on highly enriched ^{100}Mo appears to be the most promising one. Over the years several cross section measurements thereof have been reported.

However, a critical analysis of the data and the influence of co-produced long-lived Tc-isotopes on the specific activity of ^{99m}Tc need to be critically considered. From ^{100}Mo the two long-lived radioisotopes ^{99g}Tc and ^{98}Tc are co-produced. ^{99g}Tc is formed directly by the $^{100}\text{Mo}(p,2n)^{99g}\text{Tc}$ reaction and indirectly via the decay of ^{99}Mo and ^{99m}Tc during irradiation and after EOB. ^{98}Tc is generated by the $^{100}\text{Mo}(p,3n)$ reaction directly.

We calculated excitation functions for the formation of ^{99}Mo and ^{99m}Tc as well as for the long-lived technetium isotopes ^{99g}Tc and ^{98}Tc by the code TALYS for the proton-induced reactions on ^{100}Mo . For the first two nuclides calculations were also performed using the code STAPRE. The direct and indirect production of ^{99m}Tc was critically analysed. The integral yields of ^{99}Mo , ^{99m}Tc , ^{99g}Tc and ^{98}Tc were calculated for four chosen irradiation times as a function of proton energy. Therefore the activities of ^{99}Mo and ^{99m}Tc as well as the number of atoms of ^{99m}Tc , ^{99g}Tc and ^{98}Tc were deduced for six realistic proton energy ranges.

The dependence of the specific activity of ^{99m}Tc was calculated in relation to energy range, irradiation and cooling time. The specific activity of ^{99m}Tc produced directly at a cyclotron was critically compared with that obtained from a fission ^{99}Mo loaded generator. The long-lived isotopes ^{99g}Tc and ^{98}Tc cause no significant radiation and mass dose to the patient but have a strong influence on the specific activity of the cyclotron produced ^{99m}Tc depending on the production conditions. At a suggested 22 MeV incident proton energy, for example, the ratio of long-lived Tc nuclei to ^{99m}Tc nuclei may far exceed 5.0, thereby possibly affecting the kit formulation of radiopharmaceuticals and also exceeding the limits set by radiopharmaceutical regulations, e.g. in Italy. Thus, detailed experimental and theoretical investigations related to the effect of a decreasing specific activity of ^{99m}Tc on the preparation of radiopharmaceuticals appear absolutely necessary.

Primary author: SCHOLTEN, Bernhard (Forschungszentrum Jülich GmbH, Germany)

Co-authors: KONING, Arjan J. (Nuclear Research and Consultancy Group, The Netherlands); COENEN, Heinz H. (Forschungszentrum Jülich GmbH, Germany); SUDAR, Sandor (Debrecen University, Hungary); QAIM, Syed M. (Forschungszentrum Jülich GmbH, Germany)

Presenter: SCHOLTEN, Bernhard (Forschungszentrum Jülich GmbH, Germany)

Session Classification: Poster Session - Production and Application of Radionuclides

Track Classification: Production and Application of Radionuclides