



Contribution ID: 281

Type: Poster

## Measurement of the First Ionization Potentials of Short-lived Lanthanides Using a Surface Ionization Method

Tuesday, 13 May 2014 17:15 (1h 30m)

The first ionization potential (IP) directly reflects a valence electronic configuration influenced by relativistic effects which are significantly noticeable for heavy elements. Information on IP of heavy elements, therefore, gives us a better understanding of relativistic effects. IPs of heavy actinides with atomic number  $Z > 100$ , however, have not been measured by conventional techniques, such as resonance ionization mass spectrometry (RIMS), because these elements are only available in non-weighable quantities down to the one-atom-at-a-time scale. This condition forces us to call for a different experimental approach.

Here, we have focused on a surface ionization process. Since the surface ionization process takes place between an atom and a metal surface, this method is applicable to one-atom-at-a-time scale experiments. In fact, we ionized and mass-separated short-lived isotopes using the surface ionization method installed in the JAEA-ISOL system [1]. Further improvement of the surface ionization type ion-source has been recently applied to measure the IP of lawrencium (Lr,  $Z = 103$ ). 27-s  $^{256}\text{Lr}$  produced in the  $^{249}\text{Cf} + ^{11}\text{B}$  reaction was successfully ionized [2]. The IP measurement of Lr is now under way.

In this work, we measured ionization efficiencies of various short-lived lanthanides and evaluated their IPs, as a test experiment for measurement of IPs of heavy actinides.

Short-lived lanthanides,  $^{143}\text{mSm}$ ,  $^{143}\text{Eu}$ ,  $^{148}\text{mTb}$ ,  $^{154}\text{Ho}$ ,  $^{157}\text{Er}$ , and  $^{165}\text{Yb}$ , were produced by the irradiation of a 67.9-MeV  $^{11}\text{B}^{4+}$  beam delivered from the JAEA tandem accelerator on  $^{136}\text{Ce}$  /  $^{141}\text{Pr}$  /  $^{159}\text{Tb}$  and  $^{142}\text{Nd}$  /  $^{147}\text{Sm}$  /  $^{nat}\text{Eu}$  targets. Short-lived  $^{168}\text{Lu}$  was also produced in the reaction of  $^{162}\text{Dy}$  with a  $^{11}\text{B}^{4+}$  beam. Nuclear reaction products recoiling from the targets were transported to the ion-source of the JAEA-ISOL set up by a He/CdI<sub>2</sub> gas-jet transport system. The products were ionized in the ion-source, accelerated with 30 kV, mass-separated, and collected on an aluminized Mylar tape. The amounts of the collected ions were determined by gamma-ray measurement with a HP-Ge detector. To calculate ionization efficiencies, the amounts of the transported products were also determined by direct collection using a separate catcher system.

IPs of various short-lived lanthanides are evaluated based on the following Saha-Langmuir equation:

$$\alpha = n_i/n_0 = \exp((\phi - \text{IP})/kT),$$

where  $n_i$  and  $n_0$  indicate the number of ions and that of atoms on a metal surface, respectively.  $T$ ,  $k$ , and  $\phi$  are the absolute temperature of the metal surface, the Boltzmann constant, and the work function of the specific metal surface, respectively. The experimental ionization efficiency  $\beta$  is expressed by using the  $\alpha$  as follows:

$$\beta = n_i/(n_0 + n_i) = \alpha/(\alpha + 1).$$

Obtained IP values of short-lived lanthanides with tracer scale atoms were compared with literature values measured with macro-scale amounts of these elements.

[1] S. Ichikawa, et al., Nucl. Instr. and Meth. A 374, 330 (1996).

[2] T. K. Sato, et al., Rev. Sci. Instrum. 84, 023304 (2013).

**Primary author:** Mr KANEYA, Yusuke (Graduate School of Science and Engineering, Ibaraki University / Japan Atomic Energy Agency)

**Co-authors:** Dr MASATO, Asai (Japan Atomic Energy Agency); Dr SHINICHI, Ichikawa (Nishina Center for Accelerator Based Science, RIKEN); Dr SUNAO, Miyashita (Graduate school of Science, Hiroshima University); Dr

YUICHIRO, Nagame (Japan Atomic Energy Agency); Dr KAZUHIRO, Ooe (Institute of Science and Technology, Niigata University); Dr AKIHIKO, Osa (Japan Atomic Energy Agency); Dr NOZOMI, Sato (Japan Atomic Energy Agency); Dr MATTHIAS, Schädel (Japan Atomic Energy Agency); Dr THIERRY, Stora (ISOLDE, CERN); Dr SATO, Tetsuya (Japan Atomic Energy Agency); Dr ATSUSHI, Toyoshima (Japan Atomic Energy Agency); Dr KAZUAKI, Tsukada (Japan Atomic Energy Agency); Prof. JENS, V. Kratz (Institut für Kernchemie, Universität Mainz)

**Presenter:** Mr KANEYA, Yusuke (Graduate School of Science and Engineering, Ibaraki University / Japan Atomic Energy Agency)

**Session Classification:** Poster Session - Chemistry of Actinide and Trans-actinide Elements

**Track Classification:** Chemistry of Actinide and Trans-actinide Elements