



Contribution ID: 154

Type: **Invited**

## Isotopic-ratio Studies on the Redox Chemistry and Mobility of Uranium in a Phosphogypsum Disposal Site

*Friday, 16 May 2014 08:30 (30 minutes)*

The present study aims to assess the effect of redox conditions existing within a phosphogypsum disposal site on the redox stability and mobility of uranium in the disposed material. Phosphogypsum sampling and in-situ measurements were carried out at a coastal stack in Vasiliko, Cyprus. pH, EH and solubility experiments were performed in-situ and in laboratory systems. Generally, in the open stack oxidizing conditions predominate stabilizing uranium in its hexavalent oxidation state. On the other hand, after the application of a soil/vegetative cover and in the presence of natural organic matter, anoxic conditions prevail (EH < -70 mV) enabling U(VI) reduction to U(IV). Uranium concentration measurements as well as  $^{230}\text{Th}/^{238}\text{U}$  ratios show higher uranium retention under anoxic conditions, which could be ascribed to the lower solubility and mobility of U(IV).

**Primary author:** Prof. PASHALIDIS, Ioannis (Department of Chemistry, University of Cyprus)

**Co-author:** SUKSI, Juhani (Laboratory of Radiochemistry, Department of Chemistry, University of Helsinki (Fi))

**Presenter:** Prof. PASHALIDIS, Ioannis (Department of Chemistry, University of Cyprus)

**Session Classification:** Radionuclides in the Environment, Radioecology 4

**Track Classification:** Radionuclides in the Environment, Radioecology