RadChem 2014



Contribution ID: 438

Type: Verbal

Prospective carriers of 223-Ra for targeted alpha particle therapy.

Thursday, 15 May 2014 09:40 (20 minutes)

Targeted alpha particle therapy (TAT) is very powerfull tool against cancer, since DNA double-strand breaks caused by alpha radiation induce cells death.

Main advantage of radionuclides decaying by alpha particle cascade in a short-lived chain is the deposition of high energy in very small volume (approx. 27 MeV in the case of 223-Ra).

However main disadvantage of such in vivo generator is the release of daughter hot atoms from all chemical bonds. Energy of chemical bond is just several few eV compared to recoil energy of daughter atom that typically ranges to about 100 keV.

We report here on basic aspects that have to be addressed in TAT systems and describe our experimens with prospective carriers of 223-Ra intended for bone therapy. Preparation, labelling yelds and in vitro stabilities are given.

Acknowledgement

This work was partially supported by the Ministry of Education Youth and Sports of the Czech republic, grant No.: LK21310, Technological Agency of the Czech republic grant No.: TA03010027 and Czech Technical University grant No.: SGS14/084/OHK4/1D/14.

Primary authors: Dr KOZEMPEL, Jan (FJFI CVUT); Mr VLK, Martin (FJFI CVUT)

Co-authors: Ms BAJZÍKOVÁ, Anna (FJFI CVUT); Ms MÁLKOVÁ, Eva (FJFI CVUT); Dr SANTOS-OLIVEIRA, Ralph (Universidade Estadual da Zona Oeste, Brazil)

Presenter: Dr KOZEMPEL, Jan (FJFI CVUT)

Session Classification: Radiopharmaceutical Chemistry, Labelled Compounds 2

Track Classification: Radiopharmaceutical Chemistry, Labelled Compounds